

The identification of sarcosine is founded on the following observations:

1. The colour given by ninhydrin is typical for sarcosine.
2. The spot of added sarcosine coincided with the spot given by the plant extract.
3. With *p*-nitrobenzoylchloride the spot gave a yellow colour².
4. When α -amino acids were deaminated with nitrous fumes (from NaNO_2 and dilute HCl) only spots of sarcosine (characteristic colour) and proline developed when spraying the paperchromatogram with ninhydrin³.

When the plant residues after ethanol extraction were hydrolyzed with acid, sarcosine was not to be found by the paperchromatographic method. Accordingly, the proteins did not contain sarcosine.

1. Haworth, R. D., Mac Gilliway, Peacock, D. H. *Nature* **167** (1951) 1068.
2. Plattner, Pl. A., and Nager, U. *Helv. Chim. Acta* **31** (1948) 2192, 2203.
3. Dalgliesh, C. E., Johnson, A. W., Todd, A. R., and Vining, L. C. *J. Chem. Soc.* **1950** 2946.
4. Brockmann, H., and Grubhofer, N. *Naturwissenschaften* **37** (1950) 494.

Received September 30, 1953.

Carbethoxylation of 2-Thenyl Cyanide

KURT PETTERSSON

Chemical Institute, University of Uppsala,
Uppsala, Sweden

In an earlier paper the present author has described the carbethoxylation of 2-thenyl cyanide by a general process¹. This paper seems to have been overlooked by Leonard and Simet, who about a year ago described a modified procedure². During the last three years a very similar method has been used at this laboratory for the preparation of ethyl 2-thienylecyanoacetate. The differences appear to be of some interest from the practical point of view, however, and are therefore reported here.

Brändström has described a method for the carbethoxylation of ketones, esters and nitriles, using sodium ethylate (prepared from sodium and ethyl carbonate) as condensing agent, and ethyl carbonate as solvent³. A slight modification of this method (alcohol being removed *in vacuo*) gives a good yield of ethyl 2-thienylecyanoacetate.

The above-mentioned paper¹ contains a reference to a paper by Wideqvist⁴, which perhaps gives the solution of Leonard and Simet's problems concerning the hydrolysis of the alkylated 2-thienylecyanoacetates.

Experimental: 200 ml of dry ethyl carbonate was placed in a 500 ml, three-necked, round-bottomed flask, fitted with a dropping funnel, a glycerol-sealed stirrer and a Widmer column, connected to a distillation condenser. The flask was heated in an oil bath until it boiled gently. The oil bath was removed, and 3.2 g (0.14 moles) of sodium was added to the hot, stirred solution at such a rate that refluxing occurred. When all the sodium was dissolved, the mixture was allowed to cool to about 80°, and the apparatus was evacuated to a pressure of about 300 mm Hg. 16.0 g (0.13 moles) of 2-thenyl cyanide was then added in a rapid stream to the mixture in the flask. The alcohol was removed, and when the temperature at the top of the column attained the expected boiling point of ethyl carbonate, the oil bath was removed and the flask allowed to cool. The contents were poured over a mixture of 20 ml of glacial acetic acid and ice. The organic layer was separated and the water layer extracted with ether. The combined organic layers were washed with water, dried over anhydrous sodium sulphate, and the ether removed. The residue was finally fractionated *in vacuo*. The yield of ethyl 2-thienylecyanoacetate, boiling at 115°–120°/1–2 mm, was 17.7 g (70 %).

1. Pettersson, K. *Acta Chem. Scand.* **4** (1950) 395.
2. Leonard, F., and Simet, L. *J. Am. Chem. Soc.* **74** (1952) 3218.
3. Brändström, A. *Acta Chem. Scand.* **4** (1950) 1315.
4. Wideqvist, S. *Svensk Kem. Tidskr.* **55** (1943) 125; *Arkiv Kemi, Mineral. Geol.* **26A** (1948) No. 16.

Received October 10, 1953.