Enamine Chemistry

III.* A New Carboxyethylation Method

GUSTAV SCHROLL, PER KLEMMENSEN and SVEN-OLOV LAWESSON

Department of Organic Chemistry, Chemical Institute, University of Aarhus, Aarhus C., Denmark

The reaction of β -propiolactone with amines is complex and different types of reactions are observed depending upon such factors as the solvent, the amine used and the order in which the reactants are mixed. It has now been found that an enamine I, which reacts as a nucleophile (for pertinent references, see Szmuszkovicz 2) after being heated with β -propiolactone, II, smoothly and in high yields produces the corresponding δ -ketocarboxylic acid amide, III.

This new reaction, to our knowledge without precedent, implies that a rearrangement takes place as the enamine structure, I, is transformed to the carbonyl derivative, III, in the absence of water. The scope and mechanism of the reaction are now under active study and full details shall be published later. Below, a type-example is given:

Experimental. The Reaction between 1-morpholino-1-cyclohexene and β -propiolactone. 100 g (0.6 mole) of 1-morpholine-1-cyclohexene³ and 28.8 g (0.4 mole) of II in 100 ml of chlorobenzene were refluxed for 4h (inner temperature 154–155°C). The solvent and excess of enamine were distilled off and the subsequent distillation gave the main fraction with b.p. 187–188°C/1 mm Hg; $n_D^{22}=1.5090$. Yield 78 g (83 %). (Found: C 64.88; H 8.90; N 5.64; Calc.: for $C_{13}H_{21}$ O_3N (239.31): C 65.24; H 8.85; N 5.85). Mol. wt: 239 (mass spectrometry).

The 2.4-dinitrophenylhydrazone had mp. 153°C (ethanol and o-xylene). (Found: C 54.55: H 6.09; N 16.47; Calc.: for $C_{19}H_{25}O_8N_5$ (419.43); C 54.40; H 6.01; N 16.70). Mol. wt. (Rast): 428.

Basic hydrolysis of the morpholide gave a 77 % yielf of 2-oxocyclohexanepropionic acid with m.p. 64°C. (Lit.4: m.p. 64°C). Hydrolysis of the crude reaction product gave an 81 % yield of the acid.

- 1. Zaugg, H. E. Org. Reactions 8 (1954) 305.
- Szmuszkovicz, J. Advan. Org. Chem. 4 (1963) 1.
- Hünig, S., Lücke, E. and Brenninger, W. Org. Syn. 41 (1961) 65.
- Frank, R. L. and Pierle, R. C. J. Am. Chem. Soc. 73 (1951) 728.

Received November 10, 1964.

^{*} Part II: Lawesson, S.-O., Larsen, E. H. and Jakobsen, H. J. Rec. Trav. Chim. 83 (1964) 461.